



SPECIMEN LABEL

IMIDASTAR® 2L T&O

Insecticide for foliar and systemic insect control in turfgrass (including sod farms), landscape ornamentals, listed fruit and nut trees and interior plantscapes.

Manufactured for:

ALBAUGH, INC.
1525 NE 36th Street
Ankeny, Iowa 50021

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC 1-800-424-9300

For non-emergency questions for this product call Albaugh, Inc. toll free at 1-800-247-8013

AD041113A

ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]- N-nitro-2-imidazolidinimine	21.4%
OTHER INGREDIENTS	78.6%
TOTAL	100.0%

Contains 2 pounds of imidacloprid per gallon.

EPA Reg. No. 42750-115

EPA Est. No. 42750-MO-01

— SHAKE WELL BEFORE USING —

KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUCIÓN AL USUARIO: Si usted no puede leer o entender ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.
(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

In case of emergency call CHEMTREC toll free at 1-800-424-9300. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.

See inside booklet for additional PRECAUTIONARY STATEMENTS.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Keep children or pets off treated area until spray is dry.

Applicators and Other Handlers Must Wear:

1. Long-sleeved shirt and long pants
2. Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride (PVC) or viton. If you want more options, follow the instructions for category **A** on an EPA chemical-resistant category chart.
3. Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment, PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

User should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross-contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed below. In spill or leak incidents, keep unauthorized people away. You may contact Chemtrec at 1-800-424-9300 for decontamination procedures or any other assistance that may be necessary.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non-refillable containers (1, 2.5, 30 & 55 gallon): Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Offer for recycling, if available or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(non-refillable <5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(non-refillable >5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

1. Coveralls
2. Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride (PVC) or viton. If you want more options, follow the instructions for category A on an EPA chemical-resistant category chart.
3. Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets off treated area until dry.

RESTRICTION: Do not formulate this product into other end-use products.

APPLICATION TO TURFGRASS

IMIDASTAR® 2L T&O will control the following soil-inhabiting pests and larvae found in turfgrass.

PEST	SCIENTIFIC NAME
Northern & Southern masked chafers	<i>Cyclocephala borealis</i> , <i>C. immaculata</i> , and/or <i>C. lurida</i>
Asiatic garden beetle	<i>Maladera castanea</i>
European chafer	<i>Rhizotroqus majalis</i>
Green June beetle	<i>Cotinis nitida</i>
May or June beetle	<i>Phyllophaga</i> spp.
Japanese beetle	<i>Popillia japonica</i>
Oriental beetle	<i>Anomala orientalis</i>
Billbugs	<i>Spherophorus</i> spp.
Annual bluegrass weevil	<i>Hyperodes</i> spp.
Black turfgrass atenius	<i>Ataenius spretulus</i> and <i>Aphodius</i> spp.
European crane fly	<i>Tipula paludosa</i>
Mole crickets	<i>Scapteriscus</i> spp.

IMIDASTAR® 2L T&O will suppress cutworms and chinch bugs.

IMIDASTAR® 2L T&O can be applied on turfgrass in the following sites:

Home lawns	Golf courses	Playgrounds
Business and office complexes	Airports	Athletic fields
Shopping complexes	Cemeteries	Sod farms*
Multi-family residential complexes	Parks	

IMIDASTAR® 2L T&O has adequate residual activity that applications can be made preceding the egg-laying activity of the target pests.

Best control is achieved when applications are made prior to egg hatch of the pests and when rainfall or irrigation after application will penetrate vertically in the soil column carrying the active ingredient into the zone where insects are normally located. In order to move the active ingredient through the thatch, irrigate if rainfall does not occur within 24 hours after application.

Apply 1.25 to 1.6 pints per acre (equivalent to 0.46 to 0.6 fl. oz. per 1,000 sq. ft.). Make application prior to egg hatch of grubs, billbugs, annual bluegrass weevil, and European Crane Fly to maximize control.

For chinch bugs (suppression) and mole crickets apply 1.6 pints per acre (equivalent to 0.6 fl. oz. per 1,000 sq. ft.). For suppression of chinchbugs, make application before the hatching of the first instar nymphs.

For control of mole crickets make application before or during the peak egg-hatch period. If adults or large nymphs are actively tunneling, combine applications of IMIDASTAR® 2L T&O with a remedial insecticide. Follow label instructions for other insecticides when tank-mixing.

Consult your local turf State Agricultural Experiment Station or State Extension Service Specialists for more specific information regarding timing of application.

Application Equipment for Use on Turfgrass

Dilute IMIDASTAR® 2L T&O with enough water to provide adequate volume to promote thorough distribution into the pest zone. Use only accurately calibrated equipment for application to turfgrass. Apply a uniform, coarse droplet spray, using a low pressure setting to eliminate off-target drift. Perform calibration on a regular basis to ensure that equipment is distributing product properly.

RESTRICTIONS FOR TURFGRASS USE:

- Do not exceed a total of 1.6 pints application (0.4 lb. of active ingredient) per acre per year.
- Do not mow turf or lawn area until after adequate irrigation or rainfall has occurred.
- Do not apply when infested turfgrass areas are waterlogged or soil beneath turf is saturated with water. These conditions prevent thorough and consistent distribution.
- *Do not use in Arizona unless otherwise directed by supplemental labeling.

APPLICATION TO ORNAMENTALS

IMIDASTAR® 2L T&O can be applied to ornamental plants in commercial and residential landscapes and interior plantscapes. IMIDASTAR® 2L T&O is a systemic insecticide that is transported within the plant system from the roots to upper foliage. IMIDASTAR® 2L T&O must be applied into a growing area of the plant that allows absorption of the active ingredient. Adding soluble nitrogen type fertilizers to the spray solution when appropriate can promote the uptake of the active ingredient.

Application can be made by foliar application or soil applications; including soil injection, drenches, and broadcast sprays. Foliar applications offer locally systemic activity against insect pests.

The systemic translocation of active ingredient will be slower when applied to woody plants with soil applications. This delay can take 60 days or longer depending on species and size of plant. To offset this, make applications before anticipated pest infestation.

RESTRICTION: For outdoor ornamentals, applications cannot exceed a total of 1.6 pints (0.4 lb. of active ingredient) per acre per year.

ANT MANAGEMENT PROGRAMS

Use IMIDASTAR® 2L T&O to control aphids, scale insects, mealybugs and other sucking pests on ornamentals with ant populations because it removes honeydew as a food source. To enhance control of ants, supplement with residual sprays, bait placements or other ant control tactics.

APPLICATION EQUIPMENT FOR FOLIAR APPLICATIONS

IMIDASTAR® 2L T&O mixes readily with water and may be used in many types of application equipment. Add a commercial spreader/sticker to promote coverage on hard to wet foliage such as holly, pine, or ivy.

IMIDASTAR® 2L T&O is compatible with many commonly used fungicides, miticides, liquid fertilizers, and other insecticides. If applicator has no prior experience with a particular tank mix, check physical compatibility by making a small clear jar test using correct proportions of products to be tank mixed.

RESTRICTIONS:

- Keep children and pets off treated area until dry.
- Do not apply more than 1.6 pts. (0.4 lb. of active ingredient) per acre per year.
- Not for use in commercial greenhouses, nurseries, or on grasses grown for seed, or on commercial fruit and nut trees.
- Do not graze treated areas or use clippings from treated areas for feed or forage.
- Do not apply through any irrigation system.
- Do not allow puddling of irrigation water following application.
- Do not apply IMIDASTAR® 2L T&O to areas which are waterlogged or saturated, which will not allow penetration into the root zone of the plant.
- Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient.
- For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval is required.

FOLIAR APPLICATION TO ORNAMENTAL AND NON-BEARING FRUIT & NUT TREES, SHRUBS, EVERGREENS, FLOWERS, FOLIAGE PLANTS, GROUNDCOVERS AND INTERIOR PLANTSCAPES

(Only for Industrial, Commercial Buildings, Residential and Landscaped Areas)

TO CONTROL:

Adelgids	Leafminers (including Boxwood leafminer)
Aphids	Mealybugs
Honeylocust plant bug	Planthoppers
Diptera (including Rhododendron gall midge, Honeylocust pod gall midge)	Psyllids
Froghoppers	Sawfly larvae
Galls (including Hickory stem gall)	Scales (including Lecanium, Azalea bark, Calico, Cottony camellia, Cottony maple, Cottony taxus)
Japanese beetles	Spittlebugs
Lace bugs	Thrips (suppression)
Leaf beetles (including elm and viburnum leaf beetles)	(including Flower, Pear and Pine thrips)
Leafhoppers	Treehoppers
(including glassy-winged sharpshooter)	Weevils (including white Pine and Black vine)
	Whiteflies

Apply 1.5 fl. oz. (45 mL) per 100 gals. of water. Make foliar applications before high pest populations become established. Reapply on an as-needed basis.

To control White grub larvae (such as Japanese beetle larvae, Chafers, *Phyllophaga* spp., Asiatic garden beetle, Oriental beetle), apply 0.46 to 0.6 fl. oz. (14 to 17 mL) per 1,000 sq. ft.

Make broadcast applications by mixing specified rate in enough water to uniformly cover the treated area. Do not use less than 2 gals. of water per 1,000 sq. ft. In order to move the active ingredient through the thatch, irrigate if rainfall does not occur within 24 hours after application.

Refer to use directions specific for **FLOWERS** and **GROUND COVERS** concerning additional use directions.

RESTRICTION: Do not apply more than 1.6 pts. application (0.4 lb. of active ingredient) per acre per year.

SOIL APPLICATION TO ORNAMENTAL NON-BEARING FRUIT & NUT TREES, SHRUBS, FLOWERS AND GROUNDCOVERS

**(Only For Industrial, Commercial Buildings, Residential, Landscaped Planting Areas)
(Including State, National, and Private Wooded and Forested Areas)**

TO CONTROL:

Adelgids	Leafminers
Aphids	Mealybugs
Armored scales (suppression)	Pine tip moth larvae
Black vine weevil larvae	Plant bugs
Eucalyptus longhorned borer	Psyllids
Flatheaded borers (including bronze, alder and emerald ash borers)	Roundheaded borers (including Asian longhorned beetles)
Japanese beetles	Royal palm bugs
Lace bugs	Sawfly larvae
Leaf beetles (including elm and viburnum leaf beetles)	Soft scales
Leafhoppers (including glassy-winged sharpshooter)	Thrips (suppression)
	White grub larvae
	Whiteflies

For TREES –

Use the following rates as a function of tree diameter at breast height (DBH):

Apply 0.1 to 0.4 fl. oz. per inch of trunk diameter (DBH). You may use the higher rate (0.3 to 0.4 fl. oz.) only for trees greater than 15 DBH to control the following pests:

Asian longhorned beetle	Bronze birch borer
Emerald ash borer	Alder borer
Eucalyptus longhorned borer	

Restriction: Do not apply more than 25.6 fl. oz. (0.4 lb. of active ingredient) per acre per year.

Diameter at Breast Height (D.B.H.) = is measured at 4.5 feet from the ground.

APPLICATION TECHNIQUES:

SOIL INJECTION:

Mix required dosage in sufficient water to inject an equal amount of solution in each hole. For concentrate injectors mix required dose with up to 1 gallon of water per DBH inch. Maintain a low pressure and use sufficient solution for distribution of the liquid into treatment zone. Dilution rate may vary depending on equipment used, tree size and application rate. Refer to the instructions for injector equipment being used for guidance.

(continued)

- **GRID SYSTEM:** Make applications in a grid pattern on 2.5-foot centers within the drip line of the tree.
- **CIRCLE SYSTEM:** Make applications in holes evenly spaced approximately 2 to 3 feet apart in a circle within the drip line of the tree. Larger trees may require additional application circles.
- **BASAL SYSTEM:** Make applications into holes evenly spaced around the base of the tree trunk no more than 6 to 12 inches out from the base.

SOIL DRENCH:

Uniformly apply the dosage in no less than 10 gals. of water per 1,000 sq. ft. as a drench around the base of the tree, directed to the root zone. Any plastic or other barrier that may prevent drench solution from reaching the root zone must be removed.

FOR ALL APPLICATION TECHNIQUES:

- Inject an equal amount of water and solution in each hole.
- Use low pressure and sufficient solution for thorough distribution into the treatment zone.
- Maintain soil moisture for 7 to 10 days.
- Application to trees already heavily infested with borers listed may not prevent the eventual loss of the trees.

RESTRICTIONS FOR TREES:

- Do not use less than 4 holes per tree.
- No Soil Injection Applications Allowed In Nassau or Suffolk Counties of New York.
- Do not apply more than 25.6 fl. oz. (0.4 lb. of active ingredient) per acre per year.

For SHRUBS –

Apply 0.1 to 0.2 fl. oz. (3 to 6 mL) per foot of shrub height.

APPLICATION TECHNIQUES:

SOIL INJECTION:

Apply to individual plants using dosage indicated.

SOIL DRENCH:

Uniformly apply the dosage in no less than 10 gals. of water per 1,000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.

- Mix required dosage in sufficient water to inject an equal amount of solution in each hole.
- Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone.
- Maintain soil moisture for 7 to 10 days.

RESTRICTIONS FOR SHRUBS:

- Do not use less than 4 holes per shrub.
- No Soil Injection Applications Allowed In Nassau or Suffolk Counties of New York.
- Do not apply more than 25.6 fl. oz. (0.4 lb. of active ingredient) per acre per year.

For FLOWERS and GROWDCOVERS –

Apply 0.46 to 0.6 fl. oz. (14 to 17 mL) per 1,000 sq. ft.

Apply as a broadcast treatment and incorporate into the soil before planting or apply after plants are established. If application is made to established plants, irrigate after application.

POME FRUITS

(For Residential Areas Only)

(Includes: Apple, Crabapple, Loquat, Mayhaw, Pear, Pear [Oriental], Quince)

TO CONTROL:

Aphids (except Woolly apple aphid)

Leafhoppers

(including glassy-winged sharpshooter)

Leafminer

Mealybugs

San Jose Scale

Apply 1.5 fl. oz. (45 mL) per 100 gals.

OR

6.0 fl. oz. per acre as foliar spray as needed after petal fall is complete.

- For control of rosy apple aphid, apply prior to leafrolling caused by the pest.
- For first-generation leafminer control, make first application as soon as petal fall is complete. Greatest leafminer control will result from the earliest possible application. For second and succeeding generations of leafminer, optimal control is obtained from applications made early in the adult flight against egg and early instar larvae. A second application may be required 10 days later if severe pressure continues or if generations are overlapping. A single application may result in suppression only. IMIDASTAR® 2L T&O will not control late-stage larvae.
- For San Jose Scale, time applications to the crawler stage. Treat each generation.
- For late-season (preharvest) control of leafhopper species, apply IMIDASTAR® 2L T&O while most leafhoppers are in the nymphal stage.
- For optimal control of mealybug, ensure good spray coverage of the trunk and scaffolding limbs or other resting sites of the mealybug.

RESTRICTIONS FOR RESIDENTIAL POME FRUITS:

- Do not apply more than 6.0 fl. oz. per acre in a single application. Do not make more than 4 applications per acre per year.
- Allow 10 or more days between applications.
- Allow at least 7 days between last application and harvest.
- Not for use in California for control on pears.

PECAN

(For Residential Areas Only)

TO CONTROL:

Yellow pecan aphid

Pecan spittlebug

Black margined aphid

Pecan stem phylloxera

Pecan leaf phylloxera

Apply 1.5 fl. oz. (45 ml) per 100 gals.

OR

6.0 fl. oz./A¹ as a foliar application as pest pressure begins to increase.

Make a second application 10 to 14 days after first if field scouting reveals continued pest pressure. Use of an organosilicone-based spray adjuvant at specified rate can ensure thorough coverage of foliage.

(continued)

RESTRICTIONS FOR RESIDENTIAL PECAN TREES:

- Do not apply more than a total of 18.0 fl. oz. of IMIDASTAR® 2L T&O per acre per year.
- Do not make more than 3 applications per year.
- Allow 10 or more days between applications.
- Do not use on pecans in California unless directed by State supplemental labeling.

¹The amount of IMIDASTAR® 2L T&O required per acre may vary and depends on tree size or volume of foliage. The rates given are based on a standard of 400 gals. of dilute spray solutions per acre for large trees with full foliage. **DO NOT** apply more than 18 fl. oz. of IMIDASTAR® 2L T&O per acre per year.

GRAPES

(For Use Only For Industrial, Commercial Buildings and Residential Planting Areas)

TO CONTROL:

Leafhoppers (including glassy-winged sharpshooter)
Mealybugs

Apply 1.5 fl. oz. (45 mL) per 100 gals.

OR

3.0 fl. oz./A (90 mL/A) as a foliar spray using 200 gals. of water per acre.

RESTRICTIONS FOR RESIDENTIAL GRAPES:

- Do not apply more than a total of 6.0 oz. of IMIDASTAR® 2L T&O per acre per year.
- Allow at least 14 days between applications.
- Applications may be applied up to and including day of harvest.

APPLICATION TO GRASSY AREAS IN NURSERIES

This product can be used for the control of listed soil-inhabiting pests of grassy areas of nurseries castanea European chafer including Northern and Southern masked chafers, *Cyclocephala boreahs*, *C. immaculata* and/or *C. lurida*, Asiatic garden beetle, Maladera, *Rhizotroqus majalis*, Green June beetle, *Cotinis nitida*, May or June beetle, *Phyllophaga* spp., Japanese beetle, *Popillia japonica*, Oriental beetle, *Anomala orientalis*, Billbugs, *Spherophorus* spp., Annual bluegrass weevil, *Hyperodes* spp., Black turfgrass ataenius, *Ataenius spretulus* and *Aphodius* spp. and mole crickets, *Scaptenscus* spp. This product can also be used for suppression of cutworms and chinchbugs. This product can be used as directed on nursery grass in sites such as under or around field or container grown plants on roadways or other grassy areas in or around nurseries.

The active ingredient in this product has sufficient residual activity so that applications can be made preceding the egg-laying activity of the target pests. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. When applications are made prior to egg hatch of the target pests, sufficient irrigation or rainfall is needed to allow the movement of the active ingredient through the thatch.

RESTRICTIONS:

- **DO NOT** make application when grassy areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist.
- The treated grassy area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile.
- Application cannot exceed a total of 25.6 fl. oz. (1.6 pts.) (0.4 lb. of active ingredient) per acre per year.

Application Equipment for Use on Grassy Areas in Nurseries

Apply this product in sufficient water to provide adequate distribution in the treated area. The use of accurately calibrated equipment normally used for the application of soil insecticides is required. Use equipment which will produce a uniform course droplet spray using a low pressure setting to eliminate off-target drift. Check calibration periodically to ensure that equipment is working properly.

GRASSY AREAS OF FIELD AND FOREST NURSERIES

PESTS	USE RATE
Larvae of: Annual bluegrass weevil Asiatic garden beetle Billbugs Black turfgrass ataenius Cutworms (suppression) European chafer European crane fly Green June beetle Japanese beetle Northern masked chafer Oriental beetle <i>Phyllophaga</i> spp. Southern masked chafer	19.2 to 25.6 fl. oz. per acre (1.2 to 1.6 pts. per acre) or 0.46 to 0.6 fl. oz. (13 to 17 ml) per 1,000 sq. ft.
Chinchbugs (suppression) Mole crickets	25.6 fl. oz. per acre (1.6 pts. per acre) or 0.6 fl. oz. (17 ml) per 1,000 sq. ft.

Remarks:

For control of grubs, billbugs, and annual bluegrass weevil, make application prior to egg hatch of the target pest.

Read “APPLICATION EQUIPMENT” section of this label.

For suppression of chinchbugs, make application prior to or during the hatching of the first instar nymphs. For control of mole crickets make application prior to or during the peak egg-hatch period. When adults or large nymphs are present and actively tunneling, accompany the application of this product with a remedial insecticide. Follow the most restrictive label instructions when tank mixing.

Consult your local turf State Agricultural Experiment Station or State Extension Service Specialists for more specific information regarding timing of application.

RESTRICTIONS:

- Do not apply more than 25.6 fl. oz. (1.6 pts.) (0.4 lb. of active ingredient) per acre per year.
- Irrigation or rainfall must occur within 24 hours after application to move the active ingredient through the thatch.
- Do not mow turf or lawn area until after sufficient irrigation or rainfall has occurred.

APPLICATION TO ORNAMENTALS AND VEGETABLE PLANTS

(Nurseries, Greenhouses, Interior Plantscapes)

This product is for insect control on ornamental and vegetable plants in nurseries and greenhouses and interior plantscapes. This product is a systemic product and will be translocated upward into the plant system. To assure optimum effectiveness the product must be placed where the growing portion of the target plant can absorb the active ingredient. The addition of a nitrogen containing fertilizer, where applicable into the solution, may enhance the uptake of the active ingredient. Application can be made by foliar application or soil applications including soil injection drenches and broadcast sprays.

When making soil applications to plants with woody stems, systemic activity will be delayed until the active ingredient is translocated throughout the plant. Make application prior to anticipated pest infestation.

Bark Media – Media with 30% or more bark content may confer a shorter period of protection when treated with this product.

Resistance – Some insects are known to develop resistance to insecticides after repeated use. Because the development of resistance cannot be predicted, the use of this product must conform to resistance management strategies established for the use area. Consult your local or state pest management authorities for details.

Application Equipment for Ornamentals and Vegetable Plants

Mix product with the required amount of water and apply as specified for the selected use pattern.

When making foliar applications on hard-to-wet foliage such as holly pine or ivy the addition of a spreader/sticker will improve coverage. If concentrate or mist type spray equipment is used apply an equivalent amount of product on the area sprayed as would be used in a dilute application.

This product has been found to be compatible with commonly used fungicides, miticides, liquid fertilizers, and other commonly used insecticides. The physical compatibility of this product may vary with different sources of pesticide products and local cultural practices. For any tank mixture that has not been previously tested, prepare on a small scale (pint or quart jar) using the proper proportions of pesticides and water to ensure the physical compatibility of the mixture.

**FOLIAR & BROADCAST APPLICATIONS –
ORNAMENTAL TREES (including non-bearing fruit & nut trees), SHRUBS, EVERGREENS,
FLOWERS, FOLIAGE PLANTS, GROUNDCOVERS, INTERIOR PLANTSCAPES, VEGETABLE
PLANTS* (around field grown nursery and container stock, indoor and outdoor ornamentals
[including both greenhouse and interior plantscapes] and ornamentals grown in flats,
benches or beds)**

FOLIAR APPLICATIONS

PESTS	USE RATE	REMARKS
Adelgids Aphids Japanese beetles (adults) Lace bugs Leaf beetles (including Elm and Viburnum leaf beetles) Leafhoppers (including Glassy-winged sharpshooter) Leafminers Mealy bugs Sawfly larvae Thrips (suppression) Whiteflies	1.7 fl. oz. (50 ml) per 100 gals. of water	Start treatments prior to establishment of high pest populations and reapply on an as-needed basis for resistance management purposes. DO NOT make an imidacloprid foliar application following a soil application in the same crop.

BROADCAST APPLICATIONS

PESTS	USE RATE	REMARKS
White grub larvae (such as Japanese beetle larvae, chafers, <i>Phyllophaga</i> spp., Asiatic garden beetle, Oriental beetle)	0.45 to 0.6 fl. oz. (13 to 17 ml) per 1,000 sq. ft.	Mix required amount of product in sufficient water to uniformly and accurately cover the area being treated. DO NOT use less than 2 gals. of water per 1,000 sq. ft. Irrigate thoroughly to incorporate this insecticide into the upper soil profile. Refer to specific use directions (found below) for FLOWERS and GROUND COVERS .

RESTRICTIONS:

- Only for use on vegetable plants intended for resale including Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage Chinese, Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugar-beets, Tomatillo and Tomato.
- For outdoor plants grown in-ground, applications of this product cannot exceed a total of 25.6 fl. oz. (1.6 pts.) (0.4 lb. of active ingredient) per acre per year.

SOIL APPLICATIONS – NURSERY, GREENHOUSE AND INTERIORESCAPE PLANTS		
PESTS	USE RATE	APPLICATION SITE
	TREES	
Adelgids Alder borer Aphids Armored scales (suppression) Asian longhorned beetle Black vine weevil larvae Bronze birch borer Emerald ash borer Eucalyptus longhorned borer Flatheaded borers (including Bronze birch and Alder) Japanese beetles (adult) Lace bugs Leaf beetles (including Elm and Viburnum leaf beetles) Leafhoppers (including Glassy-winged sharpshooter) Leafminers Mealybugs Pine tip moth larvae Psyllids Royal palm bugs Sawfly larvae* Soft scales Thrips (suppression) White grub larvae Whiteflies	<p>For TREES Use the following rates as a function of tree diameter at breast height (DBH):</p> <p>Apply 0.1–0.4 fl. oz. (3 to 12 mL) per inch of trunk diameter (DBH)</p> <p>You may use the higher rate (0.3–0.4 fl. oz.) only for trees >15 inches (DBH) to control: Asian longhorned beetle Emerald ash borer Eucalyptus longhorned borer Bronze birch borer and Alder borer</p> <p>RESTRICTION:</p> <ul style="list-style-type: none"> • DO NOT apply more than 25.6 fl. oz. (0.4 lb. Al) per acre per year. <p>Diameter at Breast Height (DBH) is measured at 4–5 feet from the ground.</p>	<p>SOIL INJECTION:</p> <p>GRID SYSTEM: Holes must be spaced on 2- to 5-foot centers in a grid pattern extending to the drip line of the tree.</p> <p>CIRCLE SYSTEM: Apply in holes evenly spaced in circles (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line.</p> <p>BASAL SYSTEM: Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base.</p> <p>Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. DO NOT use less than 4 holes per tree.</p> <p>NEW YORK SPECIFIC RESTRICTION:</p> <ul style="list-style-type: none"> • No Soil Injection Applications Allowed in Nassau or Suffolk Counties of New York. <p>SOIL DRENCH: Uniformly apply the dosage in no less than 10 gals. of water per 1,000 sq. ft. as a drench around the base of the tree directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.</p> <p>For Control of Specified Borers Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing pest damage and tree stress.</p> <p>Basal Soil Drench using Slow Release Irrigation Bags (such as Treegator® 15-, 20- or 50-gallon capacity). Follow manufacturer’s instructions for installation and setup of drip irrigation water bag. Fill bag to 1/4 capacity with irrigation water. Add the specified rate of this product for the tree diameter to which the bag is attached. Add remaining volume of water needed to fill bag.</p>

(continued)

SOIL APPLICATIONS – NURSERY, GREENHOUSE AND INTERIORESCAPE PLANTS (cont.)		
PESTS	USE RATE	APPLICATION SITE
Adelgids Alder borer Aphids Armored scales (suppression) Asian longhorned beetle Black vine weevil larvae Bronze birch borer Emerald ash borer Eucalyptus longhorned borer Flatheaded borers (including Bronze birch and Alder) Japanese beetles (adult) Lace bugs Leaf beetles (including Elm and Viburnum leaf beetles) Leafhoppers (including Glassy-winged sharpshooter) Leafminers Mealybugs Pine tip moth larvae Psyllids Royal palm bugs Sawfly larvae* Soft scales Thrips (suppression) White grub larvae Whiteflies	0.1 to 0.2 fl. oz. (3 to 6 ml) per foot of shrub height	SOIL INJECTION: Apply to individual plants using dosage indicated. Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. DO NOT use less than 4 holes per shrub. NEW YORK SPECIFIC RESTRICTION: <ul style="list-style-type: none"> • No Soil Injection Applications Allowed in Nassau or Suffolk Counties of New York.
		SOIL DRENCH: Uniformly apply the dosage in no less than 10 gals. of water per 1,000 sq. ft. as a drench around the base of the tree directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.
		Basal Soil Drench Using Slow Release Irrigation Bags (such as Treegator® 15-, 20-, or 50-gallon capacity). Follow manufacturer’s instructions for installation and setup of drip irrigation water bag. Fill bag to 1/4 capacity with irrigation water. Add the specified rate of this product for the shrub height to which the bag is attached. Add remaining volume of water needed to fill bag.
		FLOWERS & GROUNDCOVERS
	0.1 to 0.2 fl. oz. (3 to 6 ml) per foot of shrub height	Apply as a broadcast treatment and incorporate into the soil before planting or apply after plants are established. Irrigate immediately following application to established plants.
Remarks: *Sawfly larvae feed on mature foliage beginning in early spring. Make treatments in the fall before sawfly emergence in spring to allow adequate time for imidacloprid translocation into mature foliage.		
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 25.6 fl. oz. (1.6 pts.) (0.4 lb. AI) per acre per year. • DO NOT use less than 2 gals. of spray volume per 1,000 sq. ft. (85 GPA). 		

SOIL APPLICATIONS – FIELD AND FOREST NURSERIES

PESTS	Fl. oz./1,000 ft. of row	Fl. oz./1,000 sq. ft.
For control of: White grub larvae ¹ (such as Japanese beetle, Masked chafers, European Chafer, Oriental beetle, Asiatic garden beetle)	1.7 fl. oz. (50 ml)	0.6 fl. oz. (17 ml) 25.6 fl. oz./Acre
<p>Application methods: Apply as a uniform band on either side of the row using a band width six (6) inches wider than the actual root ball diameter to be dug. DO NOT allow bands in adjacent rows to overlap.</p>		
<p>Remarks: Mowing of the vegetation in the area to be treated to a height of 3 inches or less prior to application will improve the consistency of control. Apply May through July. Time the treatment so that rainfall or irrigation occurs within 24 hours following the application. ¹For grub control in areas of turf apply as a broadcast application using 0.45–0.60 fl. oz. (13–17 ml) per 1,000 sq. ft. (19.6–25.6 fl. oz./Acre).</p>		
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT exceed 25.6 fl. oz./acre per year (1.6 pts.) (0.4 lb. AI/A). • DO NOT use less than 2 gals. of spray volume per 1,000 sq. ft. (85 GPA). 		

EBB & FLOOD APPLICATIONS

This product may be applied through Ebb and Flood applications to Ornamental and Vegetable Plants (intended for resale only) grown in containers. To assure accurate uptake prior to treatment bring a minimum of 10 plants up to a known field capacity and allow to dry out for one or two days. Re-wet these plants to determine how much water on average each plant will absorb to bring it back at field capacity. Use the volume absorbed per plant (keeping pot sizes uniform) multiplied by the number of pots being treated. Add to this volume a required minimum to flood your smallest treatment area. This will minimize the return back to the storage tank. Reuse the returned volume with subsequent irrigation or nutrients on the same plants.

(continued)

**EBB & FLOOD APPLICATIONS –
ORNAMENTAL AND VEGETABLE PLANTS GROWN IN CONTAINERS**

PESTS	CONTAINER SIZE (inches)	Herbaceous species including vegetable plants (1 or 2 plants/pot)	Woody perennials, Herbaceous species including vegetable plants (3 or more/pot)
		ml/100 plants	
Adelgids Aphids	2	1.6	2.5
Armored scales (suppression) Fungus gnats (larvae only) ¹	3	2.5	3.7
Japanese beetles (adults) Lacebugs	4	3.3	5.0
Leaf Beetles (including Elm and Viburnum Leafhoppers/ Glassy-winged Sharpshooters)	5	4.2	6.3
Leafminers Mealybugs	6	5.0	7.7
Psyllids	7	5.9	9.1
Root mealybugs	8	6.6	10.0
Root weevil complex (such as Apopka, Black vine, Citrus root) ²	9	7.4	11.1
Soft scales ³ Thrips (suppression) ⁴	10	8.3	12.5
Whiteflies	11	9.0	14.3
White grub larvae (such as Japanese beetle, Masked chafers, European)	12	10.0	16.7

Remarks:

¹Fungus gnat larvae in the soil will be controlled by drench or incorporation. No adult Fungus Gnat control. Other foliar insect control is achieved by the uptake of this product from a healthy root system translocating the active ingredient up into the plant.

²Root Mealybug control will require a thorough drenching of containerized media. Coverage is essential for control while minimizing the amount of leachate. Rate: 1.7 fl. oz. (50 ml) in 150 gals. of water.

³Citrus Root Weevil – For use on non-bearing citrus nursery stock.

⁴Thrips suppression on foliage only. Thrips in buds and flowers will not be suppressed.

RESTRICTIONS:

- For outdoor plants grown in ground applications of this product cannot exceed a total of 25.6 fl. oz. (1.6 pts.) (0.4 lb. of active ingredient) per acre per year.
- Use only on vegetable plants intended for resale including Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo and Tomato.

DRENCH & IRRIGATION APPLICATIONS

This product may be applied through Drench and Flood applications. See instructions above.

For Application Through Irrigation Systems: Apply only to greenhouse and nursery grown ornamentals, vegetable plants (intended for resale only) and interiorscape plants using soil drenches, micro-irrigation, drip irrigation, overhead irrigation, ebb and flood irrigation or hand-held or motorized calibrated irrigation equipment.

DRENCH & IRRIGATION APPLICATIONS – ORNAMENTAL AND VEGETABLE PLANTS GROWN IN SMALL CONTAINERS, IN FLATS, ON BENCHES, OR IN BEDS			
PESTS	CONTAINERIZED PLANTS (small containers)		
	CONTAINER SIZE (inches)	Herbaceous species including vegetable plants (1 or 2 plants/pot)	Woody perennials, Herbaceous species including vegetable plants (3 or more/pot)
		# of Containers treated with 2.0 fl. oz. (60 ml)	
Adelgids	2	3000	2000
Aphids	3	2000	1350
Fungus gnats (larvae only)	4	1500	1000
Japanese beetles (adults) ¹	5	1200	800
Lacebugs	6	1000	650
Leaf Beetles (including Elm and Viburnum leaf beetles)	7	850	550
Leafhoppers (including Glassy-winged sharpshooter)	8	750	500
	9	675	450
Leafminers	10	600	400
Mealybugs	11	550	350
Psyllids	12	500	300
Root mealybugs ²	Application methods: Use sufficient volume to wet most of the potting medium without loss of liquid from the bottom of the container. Apply according to label directions. Follow application with moderate irrigation. Irrigate carefully during the next 10 days in order to prevent loss of active ingredient due to leaching.		
Root weevil complex (such as Apopka, Black vine, Citrus root weevils) ³			
Soft scales			
Thrips (suppression) ⁴			
Whiteflies	PLANTS IN FLATS, ON BENCHES, OR IN BEDS		
White grub larvae (such as Japanese beetle, Masked chafer, European chafer, Oriental beetle, Asiatic garden beetle)	0.6 fl. oz. (17 ml) per 1,000 sq. ft.		
	Application Methods: Mix required amount in sufficient water to uniformly cover the area being treated. DO NOT use less than 2 gals. of mixture per 1,000 sq. ft. Apply as a broadcast treatment and incorporate into the medium before planting or apply after plants are established. Lightly water the treated areas if application is made to established plants. Allow no leaching or runoff for 10 days after application.		

(continued)

**DRENCH & IRRIGATION APPLICATIONS – ORNAMENTAL AND VEGETABLE PLANTS
GROWN IN SMALL CONTAINERS, IN FLATS, ON BENCHES, OR IN BEDS (cont.)**

Remarks:

- ¹Fungus gnat larvae in the soil will be controlled by drench or incorporation. No adult Fungus Gnat control. Other foliar insect control is achieved by the uptake of this product from a healthy root system translocating the active ingredient up into the plant.
- ²Root Mealybug control will require a thorough drenching of containerized media. Coverage is essential for control while minimizing the amount of leachate. Rate 1.7 fl. oz. (50 ml) in 150 gals. of water.
- ³Citrus Root Weevil – For use on non-bearing citrus nursery stock.
- ⁴Thrips suppression on foliage only. Thrips in buds and flowers will not be suppressed.

RESTRICTIONS:

- For outdoor plants grown in ground applications of this product cannot exceed a total of 25.6 fl. oz. (1.6 pts.) (0.4 lb. of active ingredient) per acre per year.
- Use only on vegetable plants intended for resale including Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo and Tomato.

**DRENCH AND IRRIGATION APPLICATIONS –
ORNAMENTAL AND VEGETABLE PLANTS GROWN IN LARGE CONTAINERS**

Application Instructions: Use 1.7 fl. oz. (50 ml) of product in an appropriate amount of water to prevent leaching. 1.7 fl. oz. will treat the number of containers specified below based on container size.

PESTS	CONTAINER SIZE (gals.)	# of Containers treated with 1.7 fl. oz. (50 ml)
Adelgids Aphids Fungus gnats (larvae only) ¹ Japanese beetles (adults) Lacebugs Leaf Beetles (including Elm and Viburnum leaf beetles) Leafhoppers (including Glassy-winged sharpshooter) Leafminers Mealybugs Psyllids Root mealybugs ² Root weevil complex (such as Apopka, Black vine, Citrus root) ³ Soft scales Thrips (suppression) ⁴ Whiteflies White grub larvae (such as Japanese beetle, Masked chafer, European chafer, Oriental beetle, Asiatic garden beetle)	1	340–244
	2	280–210
	3	220–165
	5	160–110
	6	100–75
	10	60–45
	15	40–30
	20	20–15

(continued)

**DRENCH AND IRRIGATION APPLICATIONS –
ORNAMENTAL AND VEGETABLE PLANTS GROWN IN LARGE CONTAINERS (cont.)**

Application Method:

Apply in sufficient water to wet the potting medium. Make applications prior to egg hatch of the target pest. Irrigate moderately after application to move the active ingredient into the root zone.

Remarks:

¹Fungus gnat larvae in the soil will be controlled by drench or incorporation. No adult Fungus Gnat control. Other foliar insect control is achieved by the uptake of this product from a healthy root system translocating the active ingredient up into the plant.

²Root Mealybug control will require a thorough drenching of containerized media. Coverage is essential for control while minimizing the amount of leachate. Rate 1.7 fl. oz. (50 ml) in 150 gals. of water.

³Citrus Root Weevil – For use on non-bearing citrus nursery stock.

⁴Thrips suppression on foliage only. Thrips in buds and flowers will not be suppressed.

RESTRICTIONS:

- For outdoor plants grown in ground applications of this product cannot exceed a total of 25.6 fl. oz. (1.6 pts.) (0.4 lb. of active ingredient) per acre per year.
- Use only on vegetable plants intended for resale including Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo and Tomato.

RESTRICTIONS:

- **DO NOT** graze treated areas or use clippings from treated areas for feed or forage. Prevent runoff or puddling of irrigation water following application.
- **DO NOT** apply this product to soils which are waterlogged or saturated which will not allow penetration into the root zone of the plants.
- Keep people and pets off treated area until dry.

Food crops

Treated areas may be replanted with any crop specified on an imidacloprid label or with any crop for which a tolerance exists for the active ingredient.

For crops not listed on an imidacloprid label or for crops for which no tolerances for the active ingredient have been established a 12-month plant-back interval must be observed.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of ALBAUGH, INC., its Supplemental Distributors, or the Seller. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, all such risks shall be assumed by the Buyer.

ALBAUGH, INC., its Supplemental Distributors and the Seller warrant that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NEITHER ALBAUGH, INC. NOR ITS SUPPLEMENTAL DISTRIBUTORS MAKE ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR ANY PARTICULAR PURPOSE OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS.**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND THE EXCLUSIVE LIABILITY OF ALBAUGH, INC., ITS SUPPLEMENTAL DISTRIBUTORS AND THE SELLER FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, When Buyer suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), Buyer must promptly notify Seller in writing of any claims to be eligible to receive either remedy stated above. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL ALBAUGH, INC., ITS SUPPLEMENTAL DISTRIBUTORS, OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.** ALBAUGH, INC., its Supplemental Distributors, and the Seller offer this product, and the Buyer accepts it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of ALBAUGH, INC.

No employee or agent of ALBAUGH, INC., its Supplemental Distributor, or the Seller is authorized to vary or exceed the terms of this Warranty in any other manner.

NOTES

NOTES

NOTES
